

DOI: 10.21767/2171-6625.1000259

The Clinical Features in Subarachnoid Hemorrhage of Unknown Etiology

Shuichi Yamada*, Hiroyuki Nakase, Young-Soo Park, Yasushi Motoyama, Ichiro Nakagawa and Fumihiko Nishimura

Department of Neurosurgery, Nara Medical University, Kashihara, Nara, Japan

*Corresponding author: Dr. Shuichi Yamada, Department of Neurosurgery, Nara Medical University, 840 Shijo-cho, Kashihara, Nara, 634-8522, Japan, Tel: +81-744-22-3051; Fax: +81-744-29-0818; E-mail: syamada@naramed-u.ac.jp

Rec Date: May 11, 2018; Acc Date: May 24, 2018, 2018; Pub Date: May 28, 2018

Citation: Yamada S, Nakase H, Park Y, Motoyama Y, Nakagawa I, et al. (2018) The Clinical Features in Subarachnoid Hemorrhage of Unknown Etiology. J Neurol Neurosci Vol.9 No.3:259.

Abstract

Background: Subarachnoid hemorrhage (SAH) without aneurysm rupture is called SAH with unknown etiology (unknown SAH). We analyzed and report the latest study of the clinical features and prognosis of patients with unknown SAH.

Methods and Findings: The study included 21 patients who had been treated at our institution. The diagnosis of unknown SAH defined as no cause of hemorrhage like cerebral aneurysm, despite of repeated vascular examination.

Discussion: The patients' age, sex, Glasgow coma scale on admission, use of antithrombotic drugs and presence of hypertension and diabetes were not concerned with prognosis. Only dyslipidemia had correlation with poor prognosis. The distribution or laterality of hematoma on computed tomography also has no relation with prognosis.

Conclusion: Our study showed no significant correlation between any characteristics of patient or SAH and prognosis, except dyslipidemia with poor prognosis. It is difficult to determine prognostic factors, therefore, we need careful examination and adequate treatment for the patients of unknown SAH.

Keywords: Subarachnoid hemorrhage; Unknown etiology; Prognosis

Introduction

Many cases of subarachnoid hemorrhage (SAH) are caused by cerebral aneurysm rupture, but there are cases of SAH not associated with aneurysm. The latter cases are referred to as "SAH of unknown etiology (unknown SAH)". It is defined as SAH without traumatic history and any vascular lesions such as aneurysm that can cause bleeding despite repeated vascular examination [1-3]. Although cases of unknown SAH have been reported in the past, the reports are often relatively old and/or

small in scale [4]. Here, we report the latest study on the clinical features and prognosis of patients with unknown SAH treated at our institution.

Research Methodology

The study included patients who had been treated for SAH at our institution from January 2007 to December 2016. Based on computed tomography (CT) scans, SAH was diagnosed in patients with grade II or greater according to the Fisher's classification. Of the patients diagnosed with SAH, those with no cerebral aneurysm, cerebral arteriovenous malformation or moyamoya disease as a likely cause of hemorrhage were diagnosed as having unknown SAH. The patients who had traumatic episode were excluded. Vascular examination was performed using either or both CT angiography and digital subtraction angiography (DSA) on at least two occasions. The imaging data were assessed by two experienced neurosurgeons and two experienced neuroradiologists. This study was pursued in accordance with the institutional review board (IRB) of our institution.

Information on patient's background characteristics, such as age, sex, Glasgow coma scale (GCS) on admission, use of antithrombotic drugs, and presence of hypertension, dyslipidemia, and diabetes as comorbidities, was collected. The type of hematoma on CT at diagnosis was classified as either local or diffused (Figure 1).




Figure 1 (A and B): Diffuse-type hematoma, with SAH at the basal cisterns and Sylvian fissures. Arrowheads show the hematoma.

Hematoma with SAH primarily localized in either the basal cistern or the Sylvian fissure was defined as local type. Hematoma with SAH found across multiple cisterns was defined as diffused type. Hematoma with SAH localized on either side of the ambient cistern or the Sylvian fissure was designated as having a left-to-right difference (laterality). Conservative therapy was administered as the initial treatment in all patients. Surgical intervention was performed for secondary hydrocephalus. Vasospasm was treated with nicardipine dosed either via drip infusion or oral route, but the medication was not used in patients with lower hematoma volume and mild symptom. Prognosis was evaluated by modified Rankin Scale (mRS) at discharge.

Statistical analysis

For univariate analysis, the chi-squared test was used to test independence from sex, GCS on admission, use of antithrombotic drugs, location of hematoma, and presence of hypertension, dyslipidemia, or diabetes. The t-test was used to analyze age (variables showing a normal distribution), whereas the Mann-Whitney U test analyzed GCS on admission (variables not showing a normal distribution). For all analyses, we used the PASW Statistics software package (version 18.0.0, SPSS Japan Inc., Tokyo, Japan). A p-value <0.05 was considered statistically significant.

Results

A total of 318 patients were studied during the defined period. Among these, 21 patients (6.6%) were diagnosed with unknown SAH and further investigated. The patient characteristics are summarized in **Table 1**.

Table 1 Characteristics of the cases.

Age (years)	63.7 (39-81)
Sex	
Male	13 (61.9)
Female	8 (38.1)
GCS on admission	14.2 (9-15)
mRS on discharge	
Favorable (0-2)	18 (85.7)
Poor (3-5)	3 (14.3)
Hypertension	
Yes	12 (57.1)
No	9 (42.9)
Dyslipidemia	
Yes	4 (19.0)
No	17 (81.0)
Diabetes	
Yes	5 (23.8)

No	16 (76.2)
Hematoma spread	
Diffuse	6 (28.6)
Local	15 (71.4)
Hematoma Laterality	
Yes	10 (47.6)
No	11 (52.4)
Antithrombotic drugs	
Yes	3 (14.3)
No	18 (85.7)
Values are presented as average (range) or n (%). GCS: Glasgow coma scale; mRS: modified Rankin scale.	

At discharge, 18 (85.7%) patients had mRS scores ranging from 0 to 2, indicating generally favorable prognosis. No patients died during the study period. Surgical intervention was used in four patients, including two patients with acute hydrocephalus treated with external ventricular drainage, one patient with delayed hydrocephalus treated with ventriculoperitoneal shunt, and one patient with adhesive arachnoiditis in the thoracic vertebra who underwent surgical dissection. Ventriculoperitoneal shunt and intervention for adhesive arachnoiditis were performed on days 33 and 450, respectively. The patients were stratified into subgroups of favorable prognosis vs. poor prognosis (mRS score, 3-5) and correlation analysis was performed on the age, sex, GCS on admission, underlying diseases, and the use of antithrombotic drugs (**Table 2**).

Table 2 Case characteristics and prognosis.

Variables	All	Prognosis		p Value
		Favorable	Poor	
Number of cases	21	18 (85.7)	3 (14.3)	n.a.
Age (years)	63.7 (39-84)	61.9 (39-81)	74.7 (59-84)	0.12
Sex (male)	13 (61.9)	11 (61.1)	2 (66.7)	0.85
GCS on admission	14.2 (9-15)	14.3 (9-15)	13.7 (12-15)	0.31
Hypertension (yes)	12 (57.1)	9 (50.0)	3 (100)	0.11
Dyslipidemia (yes)	4 (19.0)	2 (11.1)	2 (66.7)	0.02
Diabetes (yes)	5 (23.8)	3 (16.7)	2 (66.7)	0.06
Antithrombotic drugs (yes)	3 (14.3)	2 (11.1)	1 (33.3)	0.36
Values are presented as average (range) or n (%). GCS: Glasgow coma scale; n.a.: not applicable.				

The only correlation noted was a trend toward poor prognosis in patients with dyslipidemia. As for the age, the

subpopulation with poor prognosis was slightly older, but there was no significant difference. Hematoma distribution on CT was classified as either diffused or local (**Figures 1 and 2**). No significant correlation was seen between the hematoma distribution and mRS at discharge. Correlation analysis was also performed for hematoma laterality, but no significant difference was noted (**Table 3**).




Figure 2 (A and B): Local-type hematoma, with SAH localized at the basal cisterns. Arrowheads show the hematoma.

Table 3 Analysis of hematoma location and prognosis.

Variables	Prognosis		p-value
	Favorable	Poor	
Hematoma spread			0.84
Diffuse	5	1	
Local	13	2	
Hematoma laterality			0.59
Yes	9	1	
No	9	2	

Discussion

According to previous literature, the proportion of unknown SAH, that is, SAH without cerebral aneurysm/cerebral arteriovenous malformation as a likely cause of hemorrhage, is mostly between 15% and 30% [3,5-8]. However, a proportion higher than 50% has also been reported in older literature [2]. Representative reports are shown in **Table 4** [2,3,5-8]. The frequency is decreasing with the times, and this trend may be attributed to the advancement in diagnostic modalities that can detect cerebral aneurysms. In the past, traditional angiography has been used as the gold standard for the diagnosis of cerebral aneurysms, but the diagnostic capacity drastically improved with the introduction of DSA [9]. In the 2000s, diagnostic precision has further improved with the introduction of three-dimensional DSA and CT angiography [10]. This progress must have helped in reducing the frequency of unknown SAH. In particular, aneurysm in the internal carotid artery, or the so-called blister-like aneurysm, is often difficult to diagnose, and the diagnostic capacity for such a small aneurysm is likely to be highly relevant. Under the current situation, multiple examinations using different

modalities are important in order to diagnose true unknown SAH. Unlike traditional angiography, CT angiography is less invasive to patients and allows repeated examinations; hence, at least three examinations are recommended. However, examination with magnetic resonance imaging failed to diagnose unknown SAH [11].

Table 4 Transition of frequency for SAH of unknown etiology.

Authors	Year	Frequency (%)	Duration
Oder et al.	1991	41/72 (56.9)	1976-1985
van Calenbergh et al.	1993	62/294 (21.1)	Not listed
Zentner et al.	1996	86/510 (16.8)	Not listed
Flaherty et al.	2005	67/352 (19.0)	1998-2004
Gupta et al.	2009	59/353 (16.7)	2005-2006
Woznica et al.	2010	168/1084 (15.5)	1985-1992

Many previous reports, using CT, classified hematoma distribution as perimesencephalic (PM) and non-PM types. However, the frequencies of PM and non-PM types greatly vary depending on the report. The frequencies reported are 32.3% to 68.6% for PM type and 31.4% to 69.5% for non-PM type, with the predominant type varying from one report to another (**Table 5**) [3-7].

Table 5 Ratio of PM and non-PM types on CT findings.

Author	Year	PM type (%)	Non-PM type (%)
van Calenbergh et al.	1993	20/62 (32.3)	42/62 (67.7)
Canhão et al.	1995	36/71 (50.7)	35/71 (49.3)
Zentner et al.	1996	35/51 (68.6)	16/51 (31.4)
Flaherty et al.	2005	24/67 (35.8)	43/67 (64.2)
Gupta et al.	2009	18/59 (30.5)	41/59 (69.5)

PM: Perimesencephalic type; Non-PM: non-perimesencephalic type; CT: Computed Tomography.

The proportion of the PM type is also decreasing in recent reports, in the same manner as the frequency. However, such a large variation in proportion suggests that it cannot be an important factor in predicting the prognosis of unknown SAH. Therefore, we performed analysis using the classification of local vs. diffuse types to reflect hematoma volume, unlike the traditional classification. We also analyzed the laterality. This factor was not found to be a prognostic factor, however. Meanwhile, Woznica et al. reported the different conclusions. They concluded that hematoma volume on initial CT correlated with the condition at the first visit and the prognosis [8]. Their study included 168 patients with unknown SAH, a sample size vastly different from that of our study. This is the biggest limitation of the present study and has likely led to results different from theirs.

Similarly, our study did not detect a significant correlation between the prognosis and the patient's age, sex, or

underlying diseases in most cases. No correlation was seen with the use of antiplatelet agent either. The only correlation noted was a trend toward poor prognosis in patients with dyslipidemia. This difference cannot be confirmed to be significant because of the limited sample size.

Conclusion and Limitations

All we can infer is that no prognostic factor can be determined from these results. Although a favorable prognosis is likely assumed for patients with unknown SAH in many cases, a few cases have poor prognosis, which is difficult to predict. Accordingly, physicians should treat all patients with vigilance.

Other limitations of the present study include varied diagnostic and treatment methods due to the retrospective nature and possible bias in patient selection due to the single-center design. In order to analyze more cases, further investigation by multicenter prospective observational research is required.

This single-center study was conducted to investigate the clinical features and prognostic factors in patients with unknown SAH. The results did not identify a factor that may predict the prognosis, indicating that each patient should be treated in a careful and adequate manner.

References

1. Rinkel GJ, Van Gijn J, Wijdicks EF (1993) Subarachnoid hemorrhage without detectable aneurysm. A review of the causes. *Stroke* 24: 1403-1409.
2. Oder W, Kollegger H, Zeiler K, Dal-Bianco P, Wessely P, et al. (1991) Subarachnoid hemorrhage of unknown etiology: early prognostic factors for long-term functional capacity. *J Neurosurg* 74: 601-605.
3. Van Calenbergh F, Plets C, Goffin J, Velghe L (1993) Nonaneurysmal subarachnoid hemorrhage: Prevalence of perimesencephalic hemorrhage in a consecutive series. *Surg Neurol* 39: 320-323.
4. Canhão P, Ferro JM, Pinto AN, Melo TP, Campos JG (1995) Perimesencephalic and nonperimesencephalic subarachnoid hemorrhages with negative angiograms. *Acta Neurochir (Wien)* 132: 14-19.
5. Zentner J, Solymosi L, Lorenz M (1996) Subarachnoid hemorrhage of unknown etiology. *Neurol Res* 18: 220-226.
6. Flaherty ML, Haverbusch M, Kissela B, Kleindorfer D, Schneider A, et al. (2005) Perimesencephalic subarachnoid hemorrhage: Incidence, risk factors, and outcome. *J Stroke Cerebrovasc Dis* 14: 267-271.
7. Gupta SK, Gupta R, Khosla VK, Mohindra S, Chhabra R, et al. (2009) Nonaneurysmal nonperimesencephalic subarachnoid hemorrhage: is it a benign entity? *Surg Neurol* 71: 566-572.
8. Woznica M, Rosahl SK, Berlis A, Weyerbrock A (2010) Outcome correlates with blood distribution in subarachnoid hemorrhage of unknown origin. *Acta Neurochir* 152: 417-422.
9. Brant-Zawadzki M, Gould R, Norman D, Newton TH, Lane B (1983) Digital subtraction cerebral angiography by intraarterial injection: comparison with conventional angiography. *AJR Am J Roentgenol* 140: 347-353.
10. Westerlaan HE, Van Dijk JMC, Jansen-Vander Weide MC, De Groot JC, Groen RJM, et al. (2011) Intracranial aneurysms in patients with subarachnoid hemorrhage: Ct angiography as a primary examination tool for diagnosis-systematic review and meta-analysis. *Radiology* 258: 134-145.
11. Maslehaty H, Petridis AK, Barth H, Mehdorn HM (2011) Diagnostic value of magnetic resonance imaging in perimesencephalic and nonperimesencephalic subarachnoid hemorrhage of unknown origin. *J Neurosurg* 114: 1003-1007.